Wound Care Instructions

Surgery

- 1. Leave the dressing in place for 24 hours and do not get it wet.
- 2. Remove the dressing and gently wash the wound with mild soap and water. Dry the area with a clean towel. If there are crusts (scabs) on the wound then use cotton balls soaked with hydrogen peroxide to gently remove the crusts.
- 3. After cleaning the wound, apply antibiotic ointment (Polysporin or Bacitracin). Then cover the wound with a bandage. Cleaning the wound and applying a bandage is done once a day.
- 4. After 3-4 days, the wound can be left uncovered if there is no oozing of fluid or blood. It is still necessary to keep the wound moist with antibiotic ointment.

Biopsy/ Destruction of a Cancer

- 1. Leave the dressing in place for 24 hours and do not get it wet.
- 2. Remove the dressing and gently wash the wound with mild soap and water. Dry the area with a clean towel. Apply antibiotic ointment (Polysporin or Bacitracin).
- 3. Cover the wound with a bandage. Cleaning the wound and applying a bandage is done once a day until the wound is healed. This may take 1-3 weeks.

Drainage of a Cyst or Abscess

- 1. Leave the dressing in place for 24 hours and do not get it wet.
- 2. Remove the dressing and gently wash the wound with mild soap and water. Dry the area with a clean towel. Apply antibiotic ointment (Polysporin or Bacitracin).
- 3. Cover the wound with a bandage. Cleaning the wound and applying a bandage is done once a day until the wound is healed. This may take 1-3 weeks.
- 4. If there is a wick, it should be removed after 3-4 days.
- *If there is discomfort take extra strength Tylenol. Do not take more than the recommended dosage on the bottle. Non-steroidal anti-inflammatory medications such ibuprofen and naproxen are not recommended since they promote bleeding.
- * Notify us if:
 - A. Pain is not relieved with Tylenol.
 - B Wound is bleeding and cannot be stopped with 5-10 minutes of applying direct pressure.
 - C. Wound is infected. Signs of infection include pain, redness, swelling, and oozing of pus. A little amount of redness around the wound is normal.
 - D. There are any questions or concerns about the wound or wound care instructions.